

# PRECIPITATION SCAVENGING AND ATMOSPHERE-SURFACE EXCHANGE

Volume 2—The Semonin Volume:  
Atmosphere–Surface Exchange Processes

*Coordinated by*

**S. E. Schwartz**

Brookhaven National Laboratory

**W. G. N. Slinn**

Pacific Northwest Laboratory

● **HEMISPHERE PUBLISHING CORPORATION**

A member of the Taylor & Francis Group

Washington Philadelphia London

Proceedings of the Fifth International Conference on Precipitation Scavenging and Atmosphere Surface Exchange Processes, Richland, Washington, 15-19 July 1991

*SPONSORS:* U.S. Department of Energy, U.S. Environmental Protection Agency, and the Electric Power Research Institute.

*SUPPORTERS:* Atmospheric Environment Service of Environment Canada, Battelle Memorial Institute, U.S. National Oceanic and Atmospheric Administration, and U.S. National Science Foundation.

**PRECIPITATION SCAVENGING AND ATMOSPHERE-SURFACE EXCHANGE  
Volume 2 – The Semonin Volume: Atmosphere-Surface Exchange Processes**

Copyright © 1992 by Hemisphere Publishing Corporation. All rights reserved. Printed in the United States of America. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written permission of the publisher.

1 2 3 4 5 6 7 8 9 0 E B E B 9 8 7 6 5 4 3 2

Cover design by Michelle Fleitz.

A CIP catalog record for this book is available from the British Library.

∞ The paper in this publication meets the requirements of the ANSI Standard Z39.48-1984 (Permanence of Paper)

**Library of Congress Cataloging-in-Publication Data**

Precipitation scavenging and atmosphere-surface exchange : coordinated by  
S.E. Schwartz, W.G.N. Slinn.

p. cm.

"Proceedings of the Fifth International Conference on Precipitation Scavenging and Atmosphere-Surface Exchange Processes, Richland, Washington, 15-19 July 1991" — T. p. verso.

Conference sponsored by the U.S. Dept. of Energy, U.S. Environmental Protection Agency, and the Electric Power Research Institute.

Includes bibliographical references and index.

Contents: v. 1, The Georgii volume — v. 2, The Semonin volume —  
v. 3, The Summers volume.

1. Precipitation scavenging — Congresses. 2. Aerosols — Congresses. 3. Atmospheric physics — Congresses. 4. Air — Pollution — Meteorological aspects — Congresses.  
I. Schwartz, S.E. II. Slinn, W. George N., date III. United States. Dept. of Energy  
IV. United States. Environmental Protection Agency. V. Electric Power Research Institute.  
VI. International Conference on Precipitation Scavenging and Atmosphere-Surface  
Exchange Processes (5th : 1991 : Richland, Wash.)

QC882.4.P74 1992

628.5'3 — dc20

ISBN 1 56032-263 2

92-5173

CIP

## Advective Influences in the Amazonian Forest Terrain

Y. VISWANADHAM, R. C. DOS SANTOS, and R. GIELOW  
Instituto Nacional de Pesquisas Espaciais  
Secretaria Especial da Ciência e Tecnologia  
12201 São José dos Campos, SP, Brazil

### ABSTRACT

This study was conducted to evaluate the performance of the energy balance components at a Ducke Reserve Forest (DRF) site (2°57' S; 59°57' W) in the Amazon Basin, a region where a significant proportion of the energy consumed by evapotranspiration is supplied from advective sensible heat. Analyses suggest that the values of  $\gamma$  (i.e., the ratio of eddy diffusivity coefficients of heat  $K_H$  and water vapor  $K_W$ ) vary from -2.67 to 2.70 in the Richardson number range -0.17 to 0.28 and the negative sign of  $\gamma$  arises in the cases of countergradient heat flux. There is, however, evidence in the literature to support an assumption that  $\gamma > 1$  can exist under stable conditions.

### 1. INTRODUCTION

It is commonly accepted that humid tropical regions have a high radiation income and abundant precipitation so that the air is very humid and hot. It is easy to conclude that these climatic conditions generate exuberant tropical vegetation. It has long been recognized that the atmospheric and soil-vegetation systems are dynamically coupled through the physical processes which bring about the transport of thermal energy and water mass across the land surface. This phenomenon explains the importance of current knowledge regarding the micrometeorology of forests. A description of the characteristics of atmospheric turbulence in the surface layer over forest canopies is essential for the understanding of the micrometeorological environment of forests. Although measurements of mean flow and turbulent fluctuations had been made in this layer (e.g., Raupach and Thom, 1981; Shuttleworth et al., 1982; Moore, 1976; Viswanadham et al., 1987, 1990; and many others), the turbulence aspects including the advective conditions have not yet been studied sufficiently.

It is very difficult to calculate the advection in a tropical rain forest (Amazon forest) because some of the advective terms are not yet measurable due to the great

heterogeneity of species and tree sizes. Nevertheless, Warhaft (1976) provided expressions for the vertical heat and moisture fluxes in the stratified boundary layer using two different models (Donaldson 1972; Launder 1975) for the pressure strain term in the equations for the conservation of heat and moisture flux. The parameterization of Warhaft is only strictly applicable to homogeneous turbulence and we expect, at best, a qualitative view of the situation near ground. Warhaft predicted that the eddy transfer coefficient for sensible heat ( $K_H$ ) would be less than the eddy transfer coefficient for latent heat ( $K_W$ ) when the sensible heat flux ( $Q_H$ ) was directed downwards and the water vapor flux ( $Q_E$ ) was upwards, which are the conditions within an advective inversion (Warhaft, 1976). A second possible reason is that in turbulent transport,  $K_H$  and  $K_W$  depend on the characteristics of the whole flow and not simply on local properties. Thus, we expect  $K_H$  and  $K_W$  to depend on the respective temperature and humidity profile shapes and more importantly on the vertical turbulence statistical properties i.e., the vertical inhomogeneity, skewness and time scale of the turbulence, and also on the source distribution (Raupach, 1987 and Weil, 1990); if these are different, then differences between  $K_H$  and  $K_W$  are possible. The dependence of  $K_H$  on the turbulence properties and source location (bottom or top) in a convective boundary layer has been shown by Weil (1990). For example, transport within forest canopies is dominated by infrequent, large eddies, which bring air down from above the canopy and generates a negative vertical velocity skewness, where the temperature is similar to that within the canopy but humidity is lower. In these circumstances, simple K-theory is invalid and counter-gradient fluxes are observed (Deardorff, 1966; Denmead and Bradley, 1985; Raupach, 1987). A similar situation is possible within an advection inversion where the temperature is a maximum at some height but the humidity profile is monotonic (i.e., having little or lack of variation).

The present work describes the field experiments at a Ducke Reserve Forest (DRF) site (2°57'S, 59°57'W) in the Amazonian evergreen forest, Amazonas, Brazil. The main objective was to present energy fluxes under advective conditions and, to establish the relationship between eddy transfer coefficients from direct measurements of eddy fluxes of  $Q_H$  and  $Q_E$  together with simultaneous measurements of the profiles of temperature and humidity.

## 2. BASIC RELATIONSHIPS

The flow over a forest canopy of height  $h$  (m) exerts an influence throughout a roughness sublayer that extends considerably above height  $z$  (m) =  $h$ . At its upper limit, the roughness sublayer merges with an inertial sublayer in which the Monin-Obukhov's similarity theory can be applied. Generally, this similarity theory applies to horizontal homogeneous conditions. However, the Amazon advective site in DRF is a horizontal inhomogeneous one.

According to the constant flux layer similarity theory (Monin and Yaglom, 1977), the eddy transfer coefficients for sensible ( $K_H \text{ m}^2 \text{ s}^{-1}$ ) and latent ( $K_W \text{ m}^2 \text{ s}^{-1}$ ) heats may be written in terms of the fluxes and the dimensionless profile gradients as

$$K_H = -Q_H / [\rho C_p (\partial \theta / \partial z)] = k u_* (z-d) / \phi_H \quad (1)$$

and

$$K_W = -Q_E / [\rho \lambda (\partial q / \partial z)] = k u_* (z-d) / \phi_W \quad (2)$$

Also, the fluxes of momentum ( $\tau \text{ kg m}^{-1} \text{ s}^{-2}$ ), sensible ( $Q_H \text{ W m}^{-2}$ ) and latent ( $Q_E \text{ W m}^{-2}$ ) heats in terms of eddy covariances,

$$\tau = -\rho \overline{u'w'} \text{ or } u_* = (|\tau/\rho|)^{1/2} \quad (3)$$

$$Q_H = \rho C_p \overline{w'\theta'} \quad (4)$$

$$Q_E = \rho \lambda \overline{w'q'} \quad (5)$$

where  $\rho$  is the density of air ( $\text{kg m}^{-3}$ );  $k$  is von Kármán constant ( $= 0.4$ );  $u_*$  is the friction velocity ( $\text{m s}^{-1}$ );  $\theta$  is the mean potential temperature (K);  $q$  is the specific humidity ( $\text{kg kg}^{-1}$ );  $C_p$  is the specific heat of air at constant pressure ( $\text{J kg}^{-1} \text{ K}^{-1}$ );  $\lambda$  is the latent heat of vaporization of water ( $\text{J kg}^{-1}$ ),  $d$  is the zero-plane displacement (m);  $\phi_H$  and  $\phi_W$  are the nondimensional gradient functions for heat and water vapor, respectively and  $u'$ ,  $w'$ ,  $\theta'$  and  $q'$  are departures from means of the horizontal wind velocity, vertical wind velocity, potential temperature and specific humidity, respectively, and overbars denote time averages.

If we define Bowen ratios from fluxes ( $\beta_f$ ) and gradients ( $\beta_g$ ) as

$$\beta_f = (C_p/\lambda) (\overline{w'\theta'}/\overline{w'q'}) \quad (6)$$

and

$$\beta_g = (C_p/\lambda) (\Delta\theta/\Delta q) \quad (7)$$

where  $\Delta\theta$  and  $\Delta q$  are measured between the same vertical heights, and finite differences have been substituted for differentials, then from Equations (1) and (2)

$$\gamma = (K_H/K_W) = (\beta_f/\beta_g) \quad (8)$$

There are theoretical reasons why  $K_H$  may not always equal  $K_W$ . For example, buoyancy may act differently on the heat-carrying and vapor-carrying parcels of air, but this is only possible if these parcels are at least partially distinct (e.g. Launder, 1978).

The Richardson gradient number (Ri) is defined as

$$Ri = [(g/\bar{\theta}) (\partial\theta / \partial z)] / (\partial u / \partial z)^2 \quad (9)$$

where  $g$  is the acceleration due to gravity ( $m\ s^{-2}$ ) and  $\bar{\theta}$  is the average potential temperature of the layer under consideration. The above equations can be utilized to compute ratios of eddy transfer coefficients and Ri from simultaneous measurements of fluxes and profiles over the Amazon forest.

### 3. EXPERIMENTAL SITE AND MEASUREMENTS

The place selected for the experiment is in the tropical evergreen forests of the Amazonas Basin, Brazil. The measurements were made using a 45 m scaffolding tower at a site (2°57'S; 59°57'W), situated in the Ducke Reserve Forest (DRF), 26 km from Torquato Highway, Manaus, Amazonas. The altitude of the DRF site is 84 m above mean sea level. The DRF belongs to the Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil. Viswanadham et al. (1990) have presented details about the experimental site and measurements of various necessary parameters at the DRF tower.

### 4. RESULTS

This work describes some preliminary results of the advective conditions from the Amazonian rain forest micrometeorological experiment. Several intensive periods of data collection from micrometeorological instrumentation were planned from 1983 to the next few years. Already data have been collected in seven experiments. The third (13<sup>th</sup> July to 31<sup>st</sup> August, 1984) and fourth (15<sup>th</sup> March to 15<sup>th</sup> May, 1985) expeditions data sets were used to obtain various results during advective conditions. Stable conditions over the Amazon forest canopy usually occur during early morning, afternoon and night hours, when available energy (i.e., net radiation  $Q^*$ ) at the canopy surface is relatively low, so that sensible and latent heat fluxes are small and thus relatively difficult to measure. However, local advection, arising from a transition from a dry to a wet surface, typically causes locally stable conditions accompanied by very large fluxes. In other words, in an environment with warm air above colder air, turbulence moves warm air down the gradient to cooler air, i.e., a downward or negative flux. One criterion for the identification of advective influences is that  $Q_E/Q^*$  is consistently greater than unity on an hourly basis (Oke, 1979; Viswanadham and André, 1983).

The wind tunnel results of Raupach et al. (1980) show that over rough closed canopies, the logarithmic wind profile law applies only above  $h + 1.5\ell$ , where  $h$  (= 35 m) and  $\ell$  (= 2 m) are the height and the transverse length scale of the roughness elements, respectively. Viswanadham et al. (1990) have obtained that the value of  $h + 1.5\ell$  is about 38 m for

DRF site. So, measurements below 39 m were not used to compute the roughness length  $z_0$ , the zero plane displacement  $d$ , and gradients from the profiles. The iterative procedure of Robinson (1962) is used to determine the roughness length  $z_0$  (m) and  $d$  from 80 quasi-adiabatic wind profiles of the Amazon forest. The mean values of 3 m and 28 m are obtained for  $z_0$  and  $d$  respectively. Various hourly data sets were selected using the above criterion for advection situations. Equations (1-2) or Equations (6-8) are chosen to compute the ratio  $\gamma$  with the availability of the appropriate data sets for the layer 41.04 to 44.66 m above the surface. The stability parameter  $Ri$  for the same layer is obtained from Equation (9). The parameter  $Q^*$ , an important climatic characteristic, represents the amount of energy available to atmosphere and canopy processes (Figures 1-4).

Figure 1 presents fluxes of  $Q^*$ ,  $Q_H$ ,  $Q_E$ , and the residue term RE for a fine day in August 1984. There is a fairly constant partition of the available energy during daylight hours as latent and sensible heat, often some sustained evaporation into the early evening, and outgoing radiation of the order of 40 to 50  $W\ m^{-2}$  as the forest cools during the night. The suppression of turbulent interaction between the forest and the atmosphere at night, typified by the low energy fluxes, is a feature of the data and applies to all the turbulence variables. The data in Figure 1 do not satisfy energy balance at night. This may be due to an overestimation of positive  $Q_E$  and an underestimation of negative  $Q_H$  by Hydra (a battery-powered eddy correlation flux measuring device) during night hours. In fact, the detailed behavior of measured turbulent fluxes at night is difficult to interpret, and it is suspected that hourly measurements produced by the Hydra are subject to apparently haphazard fluctuations of the order of 10  $W\ m^{-2}$  (see Viswanadham et al., 1990 for discussion of Hydra measurements). This may lead to the apparent lack of energy balance closure at night in Figure 1. The term RE is subject to much erratic variation during the day period. It is assumed that the physical heat storage (i.e., soil plus canopy storage heat fluxes) is equal to RE after neglecting the biochemical heat storage due to plant photosynthesis. Heat storage within the forest canopy can be a significant portion of the energy budget for forest stands, and this is apparent in the figure. There is a loss of energy to storage in the morning as temperatures rise, and this condition reappears in the late afternoon. From Figure 1, it is evident that  $Q_E$  exceeds the net radiation before 0700 hours and after 1600 hours, and  $Q_H$  is directed toward the forest canopy. During this period the Bowen ratio  $\beta_f (=Q_H/Q_E)$  is negative and the ratio  $Q_E/Q^*$  is greater than unity for a period of 3 or 4 hours after 1600 hours. In nonhomogeneous terrain like the Amazonian forest, advection of hotter and/or drier air from upwind areas helps to increase  $Q_E$  rates. The advection of drier or warmer air will also be observed with an increase of vapor pressure deficit. Sensible heat  $Q_H$  has been drawn from the air and consumed in evaporation during advection conditions. The well-defined subsidence inversions (see de Bruin, 1983; Sá et al., 1988) and the absolute values of the

time rate of change of the vapor pressure deficits of the order  $0.5$  to  $2 \text{ mbar h}^{-1}$  were observed above the canopy. All these were responsible to the results (i.e.,  $Q_E > Q^*$ ) presented in Figure 1 for morning hours (i.e., before 0700 hours) and afternoon hours (i.e., 1600-2000 hours) over the Amazonian forest surface.




FIGURE 1. Diurnal variation of the instant energy balance components above the Amazonian rain forest during August 24, 1984.

The selected advective condition results are presented in Figure 2 which suggests good agreement between the eddy heat fluxes ( $Q_H + Q_E$ ) and the available energy ( $Q^*$ ) with a high correlation coefficient of  $r = 0.96$ . Agreement is typically better than 10%, which gives us confidence in the accuracy of the flux measurements.

The advective conditions over the Amazon forest were generally observed during the morning and evening hours. They were also noticed after the passage of local convective showers (i.e., before and after noon hours). In Figure 3, the Bowen ratio ( $\beta_f$ ) values as a function of stability parameter  $Ri$  for these conditions are presented. Sometimes these advective conditions were also observed in quasi-neutral (i.e.,  $-0.02 \leq Ri \leq 0.015$ ) stabilities (see Crawford, 1965). In such situations, the positive  $Q_H$  and  $Q_E$  values were recorded and consequently  $\beta_f$  is positive in



Figure 3. Negative  $\beta_f$  values merely indicate that the two fluxes have different signs. Figure 3 shows that this is common in advective conditions when the sensible heat flux is downward (negative), but evaporation continues so that  $Q_E$  is away from the canopy surface (positive). Actually, this was due to the importance of night radiation cooling which produced cloud fog above the forest vegetation and water condensation on the forest foliage. The importance of dew in humid tropical forest like the Amazon is well known. In the early morning hours, fog lifted and the water vapor pressure was higher than closer to the foliage surface where water vapor had condensed. This gave the impression that water vapor was advected to our system, giving positive values of  $Q_E$ .




FIGURE 2. Comparison of sum of sensible and latent fluxes ( $Q_H + Q_E$ ) measured above the Amazon forest by eddy correlation with the available energy ( $Q^*$ ) under advective conditions.

In Figure 4, the ratio  $\gamma$  as a function of  $Ri$  is presented for the advective situations. It shows that  $\gamma$  values vary from -2.67 to 2.70 in the stability range -0.17 to 0.28. Specifically, downward fluxes of  $Q_H$  were accompanied by negative gradients of specific humidity. So, in Figure 4 the ratio  $\gamma$  is positive in the advective conditions (see Equations 6 and 7). On the other hand, the negative  $\gamma$  is generally associated with negative sensible heat fluxes and

positive gradients of temperature and humidity. Thus, the best documented examples of counter-gradient flow are those for sensible and latent heat fluxes. The effects of these inconsistencies (i.e.,  $\pm Y$  values) on the calculation of scalar transport above the canopy from gradients can be seen in Figures 3 and 4, which compare the directly measured heat and water vapor fluxes above the forest with those which would be calculated from the gradients of temperature and humidity via Equations (1) to (8).




FIGURE 3. The Bowen ratio  $\beta_f (= Q_H/Q_E)$  as a function of stability parameter  $Ri$  for advective conditions.

The present results indicate that  $\gamma$  is greater than unity under stable conditions and also some negative values are noticed for  $\gamma$  (Figure 4). Literature on the subject of transfer coefficient relationships offers many diverse discussions. Rider (1954) noted approximate equality of the exchange coefficients regardless of stability conditions in most cases but exceptions were found in which the diffusivity for heat was much larger than that for vapor. At  $Ri = 0.004$  the values of  $\gamma$  were about 1.18 and 1.09 for his observation numbers 44 and 46, respectively. Pruitt and Aston (1964) concluded that  $\gamma$  was very nearly 1.0 in unstable conditions, about 1.2-1.3 at  $Ri=0$ , and as high as 2-3 under very stable conditions. Estimates of the diffusivity ratios have been given by Munn (1966) based on data from Project Green Glow at Hanford, Washington and from

O'Neill, Nebraska. The values were not obtained from fundamental eddy correlations but were derived from an indirect use of the energy balance equation. From these methods Munn (1966) presented that the values of  $\gamma$  during night time are 1.13 and 1.71 for Green Glow and O'Neill, respectively. Campbell (1973) concluded that over the stable range  $0 < Ri < 0.5$  the ratio  $\gamma \geq 1$ . For strongly stable case of  $Ri \approx 0.5$ ,  $K_H$  was sufficiently greater than  $K_W$  to lead to errors of greater than 10% in the calculations of Bowen ratio and the corresponding  $Q_E$ .




FIGURE 4. The eddy transfer coefficients ratio  $\gamma (=K_H/K_W)$  as a function of stability parameter  $Ri$  during advective conditions.

A number of experiments done in Nebraska with irrigated crops bounded by dry areas have led to comparisons of  $K_H$  and  $K_W$ . Results of Verma et al. (1978) based upon lysimetry, and Motha et al. (1979), from eddy covariances, gave  $\gamma$  of about 1.5. The analysis of Verma et al. may be questionable (Hicks and Everett, 1979; Verma et al., 1978). The work of Motha et al. (1979) should be more reliable since the eddy flux, obtained with a Gill propeller and fine wet and dry bulb thermometers, were measured separately and at the same height as the gradients. Under advective conditions, there are vertical divergences of the separate fluxes. Lang et al. (1983) obtained conflicting results for similar circumstances. With direct flux measurements, they concluded

that  $\gamma$  approached unity near neutrality and decreased to about 0.65 with increasing stability.

The above evidences suggest that  $\gamma = 1$  or  $\gamma < 1$ . There is, however, support for the condition that  $\gamma > 1$  under stable conditions. Viswanadham et al. (1987) presented a critical discussion about  $\gamma > 1$  under stable conditions. The present results are consistent with Raupach (1979) and Viswanadham et al. (1987). The most dramatic examples of negative values of  $\gamma$  associated with counter-gradient heat and water vapor fluxes over the real forest canopies. Even after applying corrections for the eddy correlations flux measurements,  $\gamma > 1$ , which conflicts with theoretical predictions of Warhaft (1976).

The discrepancy has yet to be explained. It may arise from a failure to account for all the terms in the Reynolds stress equations, from differences in the physical locations of the sources and sinks for momentum, heat and water vapor within the forest, or, perhaps, from differences in the modes of transfer of momentum and scalars at foliage surfaces (Denmead and Bradley, 1985). These results under advective conditions are important for mesoscale models which illustrate the links among the radiation budget, the climate and the circulation of the atmosphere. It was thus clear that equivalence of  $K_H$  and  $K_W$  under stable conditions should not be taken for granted and that relationships between transfer coefficients under an advective inversion need further investigation. Limited space prevents further discussions of our results.

Acknowledgments. The authors wish to thank Messrs L. D. A. Sá, A. O. Manzi and others (i.e., same as in Sá et al. 1988) for their cooperations to collect the various Amazon data sets. This work was partially supported by the Financiadora de Estudos e Projetos (FINEP), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Fundação Banco do Brasil (FBB).

## 5. REFERENCES

- Campbell A. P. (1973) The effects of stability on evaporation rates measured by the energy balance method. *Agric. Meteorol.* 11, 261-267.
- Crawford T. V. (1965) Moisture transfer in free and forced convection. *Quart. J. Roy. Meteorol. Soc.* 91, 18-27.
- Deardorff J. W. (1966) The countergradient heat flux in the lower atmosphere and in the laboratory. *J. Atmos. Sci.* 23, 503-506.
- de Bruin H. A. R. (1983) A model for the Priestley-Taylor parameter  $\alpha$ . *J. Clim. Appl. Meteorol.* 22, 572-578.

- Denmead O. T. and Bradley E. F. (1985) Flux-gradient relationships in a forest canopy. In Forest Atmospheric Interaction (Ed.) B. A. Hutchison and B. B. Hicks. D. Reidel Publ. Co., Dordrecht, pp. 421-442.
- Donaldson C. DuP. (1972) Calculations of turbulent shear flows for atmospheric and vortex motions. AIAA Journal 10, 4-12.
- Hicks B.B. and Everett R. G. (1979) Comments on "Turbulent exchange coefficients for sensible heat and water vapor under advective conditions". J. Appl. Meteorol. 18, 381-382.
- Lang A. R. G., McNaughton K. G., Fazu C., Bradley E. F., and Oftaki E. (1983) Inequality of eddy transfer coefficients for vertical transport of sensible and latent heats during advective inversions. Bound.-Layer Meteorol. 25, 25-41.
- Launder B. E. (1975) On the effects of a gravitational field on the turbulent transport of heat and momentum. J. Fluid Mech. 67, 569-581.
- Launder B. E. (1978) Heat and mass transport. In P. Bradshaw (ed.), Turbulence. 2nd Edition, Springer-Verlag, New York, pp. 238-287.
- Monin A. S. and Yaglom A. M. (1977) Statistical fluid mechanics of turbulence, Vol. 1. MIT Press, Cambridge, MA, USA, 769 pp.
- Moore C. J. (1976) Eddy flux measurements above a pine forest. Quart. J. Roy. Meteorol. Soc. 101, 913-918.
- Motha R. P., Verma S. B., and Rosenberg N. J. (1979) Exchange coefficients under sensible heat advection determined by eddy correlation. Agric. Meteorol. 20, 270-280.
- Munn R. E. (1966) Descriptive Micrometeorology. Academic Press, New York, 245 pp.
- Oke T. R. (1979) Advectively assisted evaporatranspiration from irrigated urban vegetation. Bound.-Layer Meteorol. 17, 167-173.
- Pruitt W. O. and Aston M. J. (1963) Atmospheric and surface factors affecting evapotranspiration. Final Report Task 3A99-27-005-08, Contract DA-36-036-SC-80334, University of California, Davis, pp. 69-105.
- Raupach M. R. (1979) Anomalies in flux-gradient relationships over forest. Bound.-Layer Meteorol. 16, 467-486.

- Raupach M. R. (1987) A lagrangian analysis of scalar transfer in vegetation canopies. *Quart. J. Roy. Meteorol. Soc.* 113, 107-120.
- Raupach M. R., Thom A. S., and Edwards I. (1980) A wind-tunnel study of temperature flow close to regularly arrayed rough surfaces. *Bound.-Layer Meteorol.* 18, 373-397.
- Raupach M. R. and A. S. Thom (1981) Turbulence in and above plant canopies. *Ann. Rev. Fluid Mech.* 13, 97-129.
- Rider N. E. (1954) Eddy diffusion of momentum, water vapor, and heat near the ground. *Phil. Trans. Roy. Soc. London, A*, 246, 481-501.
- Robinson S. M. (1962) Computing wind profile parameters. *J. Atmos. Sci.* 19, 189-190.
- Sá L. D. A., Viswanadham Y., and Manzi A. O. (1988) Energy flux partitioning over the Amazon forest. *Theor. Appl. Climatol.* 39, 1-16.
- Shuttleworth W. J., McNeil D. D., and Moore C. J. (1982) A switched continuous-wave sonic anemometer for measuring surface heat fluxes. *Bound.-Layer Meteorol.* 23, 425-448.
- Verma S. B., Rosenberg N. J., and Blad B. L. (1978) Turbulent exchange coefficients for sensible heat and water vapor under advection conditions. *J. Appl. Meteorol.* 17, 330-338.
- Viswanadham Y. and André R. G. B. (1983). Energy balance of soybeans grown in Brazil. *Arch. Met. Geoph. Biocl. Ser. B.* 33, 141-157.
- Viswanadham Y., Sá L. D. A., Silva Filho V. P., and Manzi A. O. (1987). Ratios of eddy transfer coefficients over the Amazon forest. *Forest Hydrology and Watershed Management (Proceedings of the Vancouver Symposium, Vancouver, Canada, 9-22 August, 1987) IAHS., Publ. n<sup>o</sup>.167, 365-373.*
- Viswanadham Y., Molion L. C. B., Manzi A. O., Sá L. D. A., Silva Filho V. P., André R. G. B., Nogueira J. L. B., and Santos R. C. (1990) Micrometeorological measurements in Amazon forest during GTE/ABLE 2A Mission. *J. Geophys. Res.* 95(D9), 13669-13682.
- Warhaft Z. (1976) Heat and moisture flux in the stratified boundary layer. *Quart. J. Roy. Meteorol. Soc.* 102, 703-707.
- Weil, J. C. (1990) A diagnosis of the asymetry in top-down and bottom-up diffusion using a Lagrangian stochastic model. *J. Atmos. Sci.*, 47, 501-515.

#### DISCUSSION

J. C. WEIL.  $K_H$  and  $K_W$  depend not on the temperature and humidity profiles but more importantly on the vertical turbulence statistical properties.....i.e., the vertical inhomogeneity, skewness, and time scale.....and also on the source distribution. The dependence of  $K_H$  on the turbulence properties and source location (bottom or top) in a convective boundary layer has been shown by Weil (1990) and Wyngaard and Weil (1991).

Y. VISWANADHAM. This information included in the text.

J. C. WEIL. In essence, you state that the countergradient vapor flux causes the negative  $\gamma$  values. Is this mainly because the canopy is transpiring and produces a positive humidity gradient?

Y. VISWANADHAM. Yes.