

to maximize the return from these kinds of observations, we developed new techniques for the production of synthetic  $z$ s from *ab initio* codes. The key problem is the production of chromatic high spatial resolution images in a reasonable amount of time at a large number of wavelengths. To calculate a radiance we have to account for the production of radiation from chemical species, energetic particle precipitation, and/or solar scattering. For all thick lines a simple line-of-sight integration is inadequate; in transport, which incurs the major computational burden, must be included. From these "forward" calculations we have been able to find simple relationships between the observed intensity and key physical parameters.

An example we will demonstrate the use of FUV airglow features to track the progress of a geomagnetic storm, show horizon images of auroral nightglow, and synoptic images of the aurora and aurora as viewed by GUVI from the TIMED orbit. We will also show how the airglow can be used to determine key parameters such as  $O/N_2$  abundance, total electron content, and energy and flux precipitating electrons in the auroral oval and discuss error rates for the retrieval of these parameters.

## 2A-19 1330h POSTER

### Remote Sensing of F-Region Ion Drifts and Ionospheric Temperature at Søndre Strømfjord, Greenland Using Doppler Measurements of the $O^+(^2P)$ State.

Carr, (Dept. of Physics, US Air Force Academy, CO 840-6425, 719-472-3510)  
Niciejewski and T L Killen (Space Physics Research Laboratory, University of Michigan, Ann Arbor, MI 109, 313-747-3425)

Ground-based observations of Doppler line profiles from the F-region  $O^+(^2P)$  state, made with the Fabry-Perot interferometer (FPI) at Søndre Strømfjord, Greenland were analyzed to provide measurements of the ionospheric velocity and ion temperature. The FPI line-of-sight (LOS) ion drift and temperature measurements have been compared with simultaneous incoherent scatter radar (ISR) measurements; the results from the two techniques are in good agreement. The ground-based FPI (320Å) observations have future utility as a routine monitor of upper thermospheric atomic oxygen number densities.

## 2A-20 1330h POSTER

### Thermospheric Gravity Wave Signatures on the Nocturnal Atomic Oxygen 630 nm Airglow at Low Latitude Over South America During Quiet and Disturbed Geomagnetic Conditions

J A Sobral, (PR/MCT - Instituto Nacional de Pesquisas Espaciais, Caixa Postal 515, 12201-970 - São José dos Campos, SP, Brazil)  
L Borba, M A Abdu, I S Batista, H Sawant, K Takahashi, C J Zamlutti, and Y Nakamura

Report photometric observations of seasonally dependent equatorward traveling perturbations of the atomic oxygen 630 nm airglow, over Cachoeira Paulista (CP, 22°41' S, 45°00' W, dip 28°S) which are apparently caused by thermospheric gravity waves. About 40 occurrences of those moving disturbances were observed from 9 nights of observations within the period of 10 August 1977 to 4 July 1984. Their equatorward velocity, wavelength and period were observed to be 150-300 m s<sup>-1</sup>, 10-250 km and 15 min-3 h, respectively. HF polarization records showed neither phase nor amplitude scintillations during the occurrence of those airglow perturbations which demonstrates that the perturbations are not associated to equatorial plasma depletions. E-SN events appear during the postsunset period and mainly within the airglow NS scan from ten minutes to few hours. It is shown that they are not necessarily related to geomagnetically active periods. A series of physical and morphological features of those phenomena are presented. The present results are, to the authors' knowledge, the first report on the detection of effects of gravity waves on the F-region atomic oxygen red airglow during undisturbed geomagnetic conditions.

## SA41A CC: 106 Thurs 0830h Mesospheric Winds, Waves, and Emissions

Presiding: T F Tuan, Univ of Cincinnati

### SA41A-1 0830h Doppler-Spread Parameterization of Gravity-Wave Momentum Deposition

C O Hines (Laboratory for Atmospheres, NASA/GSFC, Greenbelt, MD 20771; 416-977-8355; e-mail: hines@windic.yorku.ca)

The circulation of the middle atmosphere is known to be strongly influenced by the deposition of momentum by gravity waves as the waves themselves dissipate. Various parameterization schemes have been proposed in the past for use in large-scale circulation models. The scheme presented here is based on the author's Doppler-spread theory of gravity-wave saturation, in which nonlinear wave-wave interactions control the rate at which the wave spectrum is eroded as the waves propagate upward. The theory is extended here to include the effects of background winds (previously ignored) under approximations suitable for the purpose, and the momentum deposition is parameterized. So too are the corresponding heating rate, and below the turbopause, eddy diffusivity. The scheme has been applied (by others) in one particular large-scale model, with early results that are described in two following papers.

### SA41A-2 0845h In-Situ Generation of Planetary Waves Near the Mesopause During Solstices

K L Chan, J G Mengel (Both at Applied Research Corp., Landover, MD 20785)  
H G Mayr, and C O Hines (Both at NASA/GSFC, Greenbelt, MD 20771)

Using a spectral model that naturally describes the planetary waves as motions with different low order zonal wave numbers, we study the waves' generation and propagation in the mesosphere. It is found that the large north-south temperature gradient associated with the mesospheric temperature anomaly during solstices provides a background for baroclinic instability to develop, especially near the summer pole. Planetary waves with a variety of periods are generated. The most prominent are the quasi-2-day waves associated with the zonal wave number 3, such as are in fact observed most strongly in the summer hemisphere. The quasi-5-day and 2.6 day waves are associated with the zonal wave numbers 1 and 2, respectively. A trace of waves with periods around 8 days is also indicated. The planetary waves also interact with and modulate the diurnal, semi-diurnal, and ter-diurnal tides.

### SA41A-3 0900h Correlation of Small Scale Density Fluctuations and Temperature Gradients in the Mesosphere

G Lehmacher (NRC/NASA GSFC, Greenbelt, MD 20771; 301-286-1085; lehmacher@eldyn.gsfc.nasa.gov), R A Goldberg (NASA GSFC; 301-286-8603; goldberg@nssdca.gsfc.nasa.gov), and F-J Lübken (Physikalisches Institut der Universität, D-53115 Bonn, Germany; +49-228-73-2648; luebken@pi1.physik.uni-bonn.de)

The characteristics and interactions of wave motions on different scales play an important role in determining the global thermal and dynamical structure of the mesosphere and lower thermosphere. However, there is still the need for measurements which cover large scale wave structure and small scale processes simultaneously. We have obtained joint altitude profiles of temperatures, small scale fluctuations and turbulence parameters from 14 rocket soundings using neutral air number densities measured by an ionization gauge. The data were acquired between January 1990 and October 1991 during three international campaigns at high and middle latitudes and span the region from 70 to 110 km. The derivation of total number density and temperature profiles below 90 km involved supporting measurements from inflatable falling spheres. We have found in all flights high correlation between the Brunt-Vaisälä frequency squared and relative density fluctuations for large regions without notable turbulence. This observation is in agreement with the picture of adiabatic motions of air parcels which applies both to gravity waves and turbulence in a stably stratified atmosphere. Under conditions of low or negative convective stability, narrow turbulent layers with thicknesses of about 1 km have been often observed. Based on the spectra of the first mentioned "nonturbulent" fluctuations we interpret them as small saturated gravity waves. We will give a spectral estimate of the potential energy flux contained in these waves and compare our results with other studies of gravity wave spectra.

### SA41A-4 0915h Effect of 'Modal Interference' Among Gravity Wave Modes in a Windless, Realistic Atmosphere

G Munasinghe, A K Bhattacharyya and T F Tuan (Physics Department, University, Cincinnati, OH 45221-0011)

Two methods have been used for computing guided modes in the middle atmosphere: one uses the radiation boundary condition and the other uses what was tantamount to phase shift analysis. The results using these two methods disagree very strongly with each other. Tuan & Tadic (1982) showed that most of the partially guided modes obtained should cancel so that the two different methods should give the same result; that paper, however, did not provide for any explicit calculation to show this.

In the present paper, we show explicitly how, for long period gravity waves, the only middle atmospheric guided modes are the Lamb and the  $S_2$  modes and how the enormously large number of modes computed from the first method manifest themselves in the reflection coefficient: they provide the background for the reflection coefficient without providing any guidance. We also consider the case when the period of the gravity wave is lowered and how the partially guided modes first become virtually guided and then fully guided as the period is lowered to the Brunt-period range.

### SA41A-5 0930h The Saturated-Cascade Model for Atmospheric Gravity Wave Spectra, and the Wavelength-Period (W-P) Relations

E M Dewan (Phillips Laboratory, GPO, 29 Randolph Road, Hanscom AFB, MA, 01731-3010)

The purpose of this paper is to present a model explaining all the atmospheric gravity wave spectra and, in addition, a wavelength-period constraint observed by many researchers and which will henceforth be designated the W-P Relations. The power spectral densities (PSD's) in terms of horizontal,  $k_x$ , and vertical,  $k_z$ , wave numbers, and frequency,  $\omega$ , will be obtainable from this model for horizontal and vertical velocity components, temperature, and density fluctuations, etc. as a function primarily of buoyancy frequency,  $N$ , and turbulent dissipation rate  $\epsilon$ . The W-P Relations will also be functions of these parameters; and, as a result, numerous experimental predictions will be available for the purpose of testing the model. Comparisons with available data will be shown to be in reasonable agreement with the model.

### SA41A-6 0945h Damping of Atmospheric Gravity Waves Through Interaction with Molecular Vibration

L E Wharton (NASA/Goddard Space Flight Center, Laboratory for Atmospheres, Greenbelt, MD 20771; 301-286-3486)

The damping of atmospheric gravity waves through interaction with molecular vibration has been treated using a technique equivalent to that of Hines (1977). The first vibrational state of molecular nitrogen is found to produce significant damping if the interaction with carbon dioxide is included.  $CO_2$  has a vibrational level with nearly the same energy level as the lowest vibrational level of  $N_2$  and the cross section for this approximately resonant vibration state exchange dominates the quenching of the  $N_2$  vibration despite the low density of  $CO_2$ . The effect of this  $CO_2$  quenching is to move the altitude of the maximum wave damping from the mesosphere, where the effect is small, up to the base of the thermosphere, where the damping is larger. This altitude effect is very sharp as the specific heat of the  $N_2$  vibrational state has an exponential increase as the gas temperature increases to approach the vibrational temperature. In the future, the vibrational damping of gravity waves is expected to increase as the density of  $CO_2$  increases.

Reference: Hines, C. O., Relaxational dissipation in atmospheric waves, Planet. Space Sci., 25, 1045-1074, 1977.

